

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

Table 4.10 - Results of Duplicate Surface Water Sample Analysis. Units are Bq/l.
See Appendix B for the sampling locations.

Location	[RN] ^a	2×TPU ^b	MDC ^c	RER ^d	[RN]	2×TPU	MDC	RER
²⁴¹Am					²³⁸Pu			
IDN	3.05×10 ⁻⁴	7.47×10 ⁻⁴	1.42×10 ⁻⁴	0.050	0.00×10 ⁰	0.00×10 ⁰	7.59×10 ⁻⁴	0.500
IDN Dup.	2.63×10 ⁻⁴	3.74×10 ⁻⁴	3.57×10 ⁻⁴		1.96×10 ⁻⁴	3.92×10 ⁻⁴	5.29×10 ⁻⁴	
TUT	1.22×10 ⁻⁴	5.48×10 ⁻⁴	1.14×10 ⁻³	0.495	2.13×10 ⁻⁴	3.03×10 ⁻⁴	2.88×10 ⁻⁴	0.407
TUT Dup.	4.85×10 ⁻⁴	4.88×10 ⁻⁴	3.28×10 ⁻⁴		5.62×10 ⁻⁴	8.03×10 ⁻⁴	7.62×10 ⁻⁴	
²³⁹⁺²⁴⁰Pu					²³⁴U			
IDN	4.11×10 ⁻⁴	4.18×10 ⁻⁴	2.79×10 ⁻⁴	0.983	1.21×10 ⁻²	3.00×10 ⁻³	7.99×10 ⁻⁴	0.355
IDN Dup.	0.00×10 ⁰	0.00×10 ⁰	5.29×10 ⁻⁴		1.37×10 ⁻²	3.36×10 ⁻³	3.16×10 ⁻⁴	
TUT	1.06×10 ⁻⁴	2.13×10 ⁻⁴	2.88×10 ⁻⁴	0.288	9.66×10 ⁻³	2.49×10 ⁻³	2.76×10 ⁻⁴	0.818
TUT Dup.	2.80×10 ⁻⁴	5.66×10 ⁻⁴	7.62×10 ⁻⁴		1.45×10 ⁻²	5.37×10 ⁻³	8.55×10 ⁻⁴	
²³⁵U					²³⁸U			
IDN	9.36×10 ⁻⁴	8.14×10 ⁻⁴	9.84×10 ⁻⁴	0.066	1.22×10 ⁻²	3.00×10 ⁻³	2.92×10 ⁻⁴	0.693
IDN Dup.	1.01×10 ⁻³	7.81×10 ⁻⁴	3.89×10 ⁻⁴		9.40×10 ⁻³	2.61×10 ⁻³	8.55×10 ⁻⁴	
TUT	5.03×10 ⁻⁴	5.07×10 ⁻⁴	3.40×10 ⁻⁴	0.122	7.81×10 ⁻³	2.15×10 ⁻³	2.75×10 ⁻⁴	0.704
TUT Dup.	3.89×10 ⁻⁴	7.84×10 ⁻⁴	1.05×10 ⁻³		5.33×10 ⁻³	2.86×10 ⁻³	8.51×10 ⁻⁴	

^a [RN] = Radionuclide concentration

^b Total propagated uncertainty

^c Minimum detectable concentration

^d Relative Error Ratio

4.6 Soil Samples

4.6.1 Sampling

Soil samples were collected from near the low-volume air samplers at six different locations around the WIPP site: MLR, SEC, SMR, WEE, WFF, and WSS (Figure 4.6). Samples were collected from each location in three incremental profiles: surface soil (SS, 0-2 cm [0-0.8 in.]), intermediate soil (SI, 2-5 cm [0.8-2 in.]), and deep soil (SD, 5-10 cm [2-4 in.]). Measurements of radionuclides in depth profiles provide information about their vertical movements in the soil systems.

4.6.2 Sample Preparation

Soil samples were dried at 110°C (230°F) for several hours and homogenized by grinding to small particle sizes. One gram (0.04 oz) of soil was dissolved by heating it with a mixture of nitric, hydrochloric, and hydrofluoric acids. Finally, it was heated with nitric and boric acids, and the residue was dissolved in hydrochloric acid for the determination of individual radionuclides.

4.6.3 Determination of Individual Radionuclides

Gamma-emitting radionuclides (⁴⁰K, ⁶⁰Co, and ¹³⁷Cs) were determined by counting an aliquot of well-homogenized ground soil samples by gamma-spectrometry. Strontium-90 was analyzed from an aliquot of the sample solution by separating it from other stable and radioactive elements using radiochemical techniques and beta

Waste Isolation Pilot Plant 2001 Site Environmental Report

DOE/WIPP 02-2225

counting. Another aliquot of the sample solution was used for the sequential determinations of alpha-emitting radionuclides, such as ^{234}U , ^{235}U , and ^{238}U ; ^{238}Pu and $^{239+240}\text{Pu}$; and ^{241}Am . These radionuclides were separated from the bulk of the inorganic materials present in the soil samples and from one another by radiochemical separations including co-precipitation and ion-exchange chromatography. Finally, the samples were micro-precipitated, filtered onto micro-filters, and counted on the alpha spectrometer.



Figure 4.6 - Soil and Vegetation Sampling Areas in 2001

4.6.4 Results and Discussions

Uranium-234 and ^{238}U were detected in every soil sample in 2001 and ^{235}U was detected in all but one. Concentrations of ^{234}U in surface soils (0-2 cm) ranged from a minimum of $5.66 \times 10^{-3} \pm 1.35 \times 10^{-3} \text{ Bq/g}$ ($1.53 \times 10^{-1} \pm 3.65 \times 10^{-2} \text{ pCi/g}$) at WEE to a maximum of $1.25 \times 10^{-2} \pm 2.48 \times 10^{-3} \text{ Bq/g}$ ($3.38 \times 10^{-1} \pm 6.70 \times 10^{-2} \text{ pCi/g}$) at SEC (Table 4.11). Concentrations of ^{235}U in the same samples ranged from $2.83 \times 10^{-4} \pm 2.72 \times 10^{-4} \text{ Bq/g}$ ($7.65 \times 10^{-3} \pm 7.35 \times 10^{-3} \text{ pCi/g}$) at WFF to $2.32 \times 10^{-3} \pm 2.48 \times 10^{-4} \text{ Bq/g}$ ($6.27 \times 10^{-2} \pm 6.70 \times 10^{-3} \text{ pCi/g}$) at WEE. The concentration of ^{238}U in surface soils ranged from $6.85 \times 10^{-3} \pm 1.57 \times 10^{-3} \text{ Bq/g}$ ($1.85 \times 10^{-1} \pm 4.24 \times 10^{-2} \text{ pCi/g}$) at WEE to $1.28 \times 10^{-2} \pm 2.63 \times 10^{-3} \text{ Bq/g}$ ($3.46 \times 10^{-1} \pm 7.11 \times 10^{-1} \text{ pCi/g}$) at WFF.

The results for uranium in intermediate depth (2-5 cm) soil samples are also given in Table 4.11. The concentration of ^{234}U ranged from $6.73 \times 10^{-3} \pm 1.38 \times 10^{-3} \text{ Bq/g}$ ($1.82 \times 10^{-1} \pm 3.73 \times 10^{-2} \text{ pCi/g}$) at WEE to $2.03 \times 10^{-2} \pm 3.92 \times 10^{-3} \text{ Bq/g}$ ($5.49 \times 10^{-1} \pm$

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

1.06×10⁻¹ pCi/g) at SEC. Uranium-235 in these soils was lowest at WEE (3.92×10⁻⁴ ± 2.43×10⁻⁴ Bq/g [1.06×10⁻² ± 6.57×10⁻³ pCi/g]) and highest at SEC (1.11×10⁻³ ± 4.92×10⁻⁴ Bq/g [3.00×10⁻² ± 1.16×10⁻² pCi/g]). The concentration of ²³⁸U ranged from 7.33×10⁻³ ± 1.48×10⁻³ Bq/g (1.98×10⁻¹ ± 4.00×10⁻² pCi/g) at WEE to 2.27×10⁻² ± 4.33×10⁻³ Bq/g (6.14×10⁻¹ ± 1.17×10⁻¹ pCi/g) at SEC.

Table 4.11 - Uranium Concentrations (Bq/g) in Soil Near the WIPP Site.
See Appendix B for the sampling locations.

Location	Depth (cm)	²³⁴ U			²³⁵ U			²³⁸ U		
		[RN] ^a	2 × TPU ^b	MDC ^c	[RN]	2 × TPU	MDC	[RN]	2 × TPU	MDC
MLR	0-2	1.20×10 ⁻²	2.22×10 ⁻³	2.19×10 ⁻⁴	4.77×10 ⁻⁴	2.75×10 ⁻⁴	9.95×10 ⁻⁵	1.13×10 ⁻²	2.10×10 ⁻³	8.03×10 ⁻⁵
MLR	2-5	1.41×10 ⁻²	2.45×10 ⁻³	7.22×10 ⁻⁵	4.92×10 ⁻⁴	2.66×10 ⁻⁴	8.92×10 ⁻²	1.32×10 ⁻²	2.32×10 ⁻³	2.47×10 ⁻⁴
MLR	5-10	1.18×10 ⁻²	2.13×10 ⁻³	6.99×10 ⁻⁵	6.36×10 ⁻⁴	3.00×10 ⁻⁴	8.62×10 ⁻⁵	1.17×10 ⁻²	2.11×10 ⁻³	6.96×10 ⁻⁵
SEC	0-2	1.25×10 ⁻²	2.48×10 ⁻³	2.47×10 ⁻⁴	1.03×10 ⁻³	4.48×10 ⁻⁴	1.12×10 ⁻⁴	1.28×10 ⁻²	2.52×10 ⁻³	9.03×10 ⁻⁵
SEC	2-5	2.03×10 ⁻²	3.92×10 ⁻³	2.77×10 ⁻⁴	1.11×10 ⁻³	4.92×10 ⁻⁴	1.25×10 ⁻⁴	2.27×10 ⁻²	4.33×10 ⁻³	1.01×10 ⁻⁴
SEC	5-10	1.90×10 ⁻²	3.50×10 ⁻³	8.77×10 ⁻⁵	8.40×10 ⁻⁴	4.07×10 ⁻⁴	2.95×10 ⁻⁴	1.61×10 ⁻²	3.02×10 ⁻³	2.38×10 ⁻⁴
SMR	0-2	1.04×10 ⁻²	1.92×10 ⁻³	7.18×10 ⁻⁵	4.59×10 ⁻⁴	2.55×10 ⁻⁴	8.84×10 ⁻⁵	1.09×10 ⁻²	2.00×10 ⁻³	7.14×10 ⁻⁵
SMR	2-5	1.11×10 ⁻²	2.02×10 ⁻³	7.22×10 ⁻⁵	7.22×10 ⁻⁴	3.27×10 ⁻³	8.88×10 ⁻⁵	1.14×10 ⁻²	2.06×10 ⁻³	7.18×10 ⁻⁵
SMR	5-10	2.09×10 ⁻²	3.74×10 ⁻³	8.51×10 ⁻⁵	8.92×10 ⁻⁴	4.00×10 ⁻⁴	1.05×10 ⁻⁴	2.09×10 ⁻²	3.74×10 ⁻³	8.47×10 ⁻⁵
WEE	0-2	5.66×10 ⁻³	1.35×10 ⁻³	1.02×10 ⁻⁴	2.32×10 ⁻⁵	2.48×10 ⁻⁴	3.41×10 ⁻⁴	6.85×10 ⁻³	1.57×10 ⁻³	1.01×10 ⁻⁴
WEE	2-5	6.73×10 ⁻³	1.38×10 ⁻³	7.81×10 ⁻⁵	3.92×10 ⁻⁴	2.43×10 ⁻⁴	9.62×10 ⁻⁵	7.33×10 ⁻³	1.48×10 ⁻³	7.77×10 ⁻⁵
WEE	5-10	1.84×10 ⁻²	3.68×10 ⁻³	2.83×10 ⁻⁴	4.74×10 ⁻⁴	3.11×10 ⁻⁴	1.28×10 ⁻⁴	2.00×10 ⁻²	3.96×10 ⁻³	1.04×10 ⁻⁴
WFF	0-2	1.17×10 ⁻²	2.45×10 ⁻³	1.04×10 ⁻⁴	2.83×10 ⁻⁴	2.72×10 ⁻⁴	3.48×10 ⁻⁴	1.28×10 ⁻²	2.63×10 ⁻³	2.80×10 ⁻⁴
WFF	2-5	1.22×10 ⁻²	2.62×10 ⁻³	3.56×10 ⁻⁴	6.62×10 ⁻⁴	3.74×10 ⁻⁴	1.28×10 ⁻⁴	1.23×10 ⁻²	2.62×10 ⁻³	2.81×10 ⁻⁴
WFF	5-10	1.00×10 ⁻²	2.18×10 ⁻³	1.08×10 ⁻⁴	2.96×10 ⁻⁴	2.47×10 ⁻⁴	1.34×10 ⁻⁴	9.36×10 ⁻³	2.05×10 ⁻³	1.08×10 ⁻⁴
WSS	0-2	1.24×10 ⁻²	2.37×10 ⁻³	2.92×10 ⁻⁴	4.26×10 ⁻⁴	2.66×10 ⁻⁴	1.05×10 ⁻⁴	1.25×10 ⁻²	2.39×10 ⁻³	2.30×10 ⁻⁴
WSS	2-5	9.58×10 ⁻³	1.97×10 ⁻³	9.25×10 ⁻⁵	7.99×10 ⁻⁴	3.92×10 ⁻⁴	1.14×10 ⁻⁴	9.18×10 ⁻³	1.53×10 ⁻³	9.18×10 ⁻⁵
WSS	5-10	8.73×10 ⁻³	1.86×10 ⁻³	9.77×10 ⁻⁵	1.78×10 ⁻⁴	1.81×10 ⁻⁴	1.21×10 ⁻⁴	8.51×10 ⁻³	1.82×10 ⁻³	9.73×10 ⁻⁵

^a [RN] = Radionuclide concentration

^b Total propagated uncertainty

^c Minimum detectable concentration

Concentrations of ²³⁴U, ²³⁵U, and ²³⁸U were also measured in deep soils (5-10 cm) (Table 4.11). Concentrations of ²³⁴U varied from 8.73×10⁻³ ± 1.86×10⁻³ Bq/g (2.36×10⁻¹ ± 5.03×10⁻² pCi/g) at WSS to 2.09×10⁻² ± 3.74×10⁻³ Bq/g (5.65×10⁻¹ ± 1.01×10⁻¹ pCi/g) at SMR. The lowest concentration of ²³⁵U in deep soils was found at WSS (1.78×10⁻⁴ ± 1.81×10⁻⁴ Bq/g [4.81×10⁻³ ± 4.89×10⁻³ pCi/g]) and the highest concentration was found at SMR (8.92×10⁻⁴ ± 4.00×10⁻⁴ Bq/g [2.41×10⁻² ± 1.08×10⁻² pCi/g]). Uranium-238 lowest concentration was 8.51×10⁻³ ± 1.82×10⁻³ Bq/g (2.30×10⁻¹ ± 4.92×10⁻² pCi/g) at WSS and the highest was found 2.09×10⁻² ± 3.74×10⁻³ Bq/g (5.65×10⁻¹ ± 1.01×10⁻¹ pCi/g) at SMR.

No uranium isotope varied significantly with year 2001 (ANOVA, ²³⁴U p = 0.105, ²³⁸U p = 0.722). All maximum measured concentrations fell within the range of natural concentrations of uranium found in soils throughout the world (Pais and Jones 1997). All these results suggest a pattern of natural variability consistent with the existence of natural uranium, without amendment from artificial sources.

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

Plutonium-238, $^{239+240}\text{Pu}$, and ^{241}Am were also analyzed in these soil samples (Table 4.12). Plutonium-238 was not detected in any samples. The measured concentration of $^{239+240}\text{Pu}$ was greater than the MDC at location MLR at all three depths. The ^{241}Am concentration was greater than the MDC at MLR, 0-2 cm and 2-5 cm, SEC, 5-10cm, and WEE, 5-10cm. Historically, soil samples collected in the same locations have shown positive results on numerous occasions. Since 1997, soil samples collected by the Environmental Monitoring group at WEE, SEC, and MLR have shown levels of ^{241}Am and $^{239+240}\text{Pu}$ slightly above background. During this time period, three different analytical laboratories were used; all had similar results. The source of activity in WIPP samples could be due to natural transport of contaminated soil from the Gnome Site via wind. The Gnome Site lies about 9km southwest of the WIPP boundary and was contaminated with fission products in 1961 when an underground test of a 3-kiloton ^{239}Pu device vented to the surface. Because there are elevated levels of radionuclides in the soil near the Gnome Site, there is potential for contamination of WIPP environmental samples. The measurable levels of radionuclides remains relatively high around the Gnome Site despite remediation efforts.

Table 4.12 - Americium and Plutonium Concentrations (Bq/g) in Soil Near the WIPP Site.
See Appendix B for the sampling locations.

Location	Depth (cm)	^{241}Am			^{238}Pu			$^{239+240}\text{Pu}$		
		[RN] ^a	2 × TPU ^b	MDC ^c	[RN]	2 × TPU	MDC	[RN]	2 × TPU	MDC
MLR	0-2	1.79×10 ⁻⁴	1.49×10 ⁻⁴	8.07×10 ⁻⁵	-3.43×10 ⁻⁵	6.88×10 ⁻⁵	2.52×10 ⁻⁴	3.42×10 ⁻⁴	2.23×10 ⁻⁴	9.29×10 ⁻⁵
MLR	2-5	4.18×10 ⁻⁴	2.29×10 ⁻⁴	8.10×10 ⁻⁵	0.00×10 ⁰	0.00×10 ⁰	1.28×10 ⁻⁴	6.59×10 ⁻⁴	3.70×10 ⁻⁴	1.28×10 ⁻⁴
MLR	5-10	2.41×10 ⁻⁴	2.31×10 ⁻⁴	3.19×10 ⁻⁴	-4.88×10 ⁻⁵	9.84×10 ⁻⁵	3.60×10 ⁻⁴	5.85×10 ⁻⁴	3.55×10 ⁻⁴	1.32×10 ⁻⁴
SEC	0-2	3.23×10 ⁻⁵	1.45×10 ⁻⁴	3.00×10 ⁻⁴	4.44×10 ⁻⁵	8.92×10 ⁻⁵	1.20×10 ⁻⁴	1.77×10 ⁻⁴	1.80×10 ⁻⁴	1.20×10 ⁻⁴
SEC	2-5	8.95×10 ⁻⁵	1.34×10 ⁻⁴	2.20×10 ⁻⁴	1.70×10 ⁻⁴	2.10×10 ⁻⁴	3.13×10 ⁻⁴	1.70×10 ⁻⁴	1.72×10 ⁻⁴	1.15×10 ⁻⁴
SEC	5-10	1.37×10 ⁻⁴	1.16×10 ⁻⁴	8.99×10 ⁻⁴	6.29×10 ⁻⁵	1.26×10 ⁻⁴	2.31×10 ⁻⁴	6.25×10 ⁻⁵	1.26×10 ⁻⁴	2.31×10 ⁻⁴
SMR	0-2	0.00×10 ⁰	0.00×10 ⁰	2.21×10 ⁻⁴	-4.07×10 ⁻⁵	8.14×10 ⁻⁵	2.99×10 ⁻⁴	0.00×10 ⁰	0.00×10 ⁰	2.99×10 ⁻⁴
SMR	2-5	8.03×10 ⁻⁵	1.15×10 ⁻⁴	1.09×10 ⁻⁴	0.00×10 ⁰	0.00×10 ⁰	1.30×10 ⁻⁴	0.00×10 ⁰	0.00×10 ⁰	3.52×10 ⁻⁴
SMR	5-10	2.98×10 ⁻⁵	1.34×10 ⁻⁴	2.78×10 ⁻⁴	-6.62×10 ⁻⁵	9.44×10 ⁻⁵	3.07×10 ⁻⁴	3.30×10 ⁻⁵	1.15×10 ⁻⁴	2.43×10 ⁻⁴
WEE	0-2	6.33×10 ⁻⁵	9.03×10 ⁻⁵	8.58×10 ⁻⁵	0.00×10 ⁰	0.00×10 ⁰	9.81×10 ⁻⁵	3.61×10 ⁻⁵	7.25×10 ⁻⁵	9.81×10 ⁻⁵
WEE	2-5	0.00×10 ⁰	0.00×10 ⁰	3.47×10 ⁻⁴	3.96×10 ⁻⁵	7.92×10 ⁻⁵	1.07×10 ⁻⁴	1.58×10 ⁻⁴	1.60×10 ⁻⁴	1.07×10 ⁻⁴
WEE	5-10	1.32×10 ⁻⁴	1.54×10 ⁻⁴	1.19×10 ⁻⁴	0.00×10 ⁰	0.00×10 ⁰	2.25×10 ⁻⁴	9.18×10 ⁻⁵	1.07×10 ⁻⁴	8.29×10 ⁻⁵
WFF	0-2	1.68×10 ⁻⁴	1.71×10 ⁻⁴	1.14×10 ⁻⁴	4.14×10 ⁻⁵	8.29×10 ⁻⁵	1.12×10 ⁻⁴	1.65×10 ⁻⁴	1.67×10 ⁻⁴	1.12×10 ⁻⁴
WFF	2-5	1.96×10 ⁻⁴	2.09×10 ⁻⁴	3.03×10 ⁻⁴	8.51×10 ⁻⁵	2.95×10 ⁻⁴	6.25×10 ⁻⁴	8.51×10 ⁻⁵	1.71×10 ⁻⁴	2.30×10 ⁻⁴
WFF	5-10	5.62×10 ⁻⁵	1.95×10 ⁻⁴	4.14×10 ⁻⁴	1.18×10 ⁻⁴	1.77×10 ⁻⁴	2.89×10 ⁻⁴	7.84×10 ⁻⁵	1.12×10 ⁻⁴	1.06×10 ⁻⁴
WSS	0-2	-3.25×10 ⁻⁵	1.45×10 ⁻⁴	3.50×10 ⁻⁴	0.00×10 ⁰	0.00×10 ⁰	9.21×10 ⁻⁵	1.36×10 ⁻⁴	1.38×10 ⁻⁴	9.21×10 ⁻⁵
WSS	2-5	7.29×10 ⁻⁵	1.04×10 ⁻⁴	9.48×10 ⁻⁵	-3.53×10 ⁻⁵	1.22×10 ⁻⁴	3.27×10 ⁻⁴	7.03×10 ⁻⁵	1.00×10 ⁻⁴	9.55×10 ⁻⁵
WSS	5-10	1.21×10 ⁻⁴	1.73×10 ⁻⁴	2.82×10 ⁻⁴	9.62×10 ⁻⁵	1.37×10 ⁻⁴	1.30×10 ⁻⁴	4.81×10 ⁻⁵	1.67×10 ⁻⁴	3.54×10 ⁻⁴

^a [RN] = Radionuclide concentration

^b Total propagated uncertainty

^c Minimum detectable concentration

Potassium-40, as expected, was detected in every sample (Table 4.13). This naturally-occurring gamma-emitting radionuclide is ubiquitous in soils. Concentrations in surface soils ranged from $2.24 \times 10^{-1} \pm 3.05 \times 10^{-2}$ Bq/g ($6.05 \times 10^0 \pm 8.24 \times 10^{-1}$ pCi/g) at SEC to $4.74 \times 10^{-1} \pm 6.18 \times 10^{-2}$ Bq/g ($1.28 \times 10^1 \pm 1.67 \times 10^0$ pCi/g) at SMR. In intermediate

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

depth soils, concentrations of ^{40}K varied from $2.32 \times 10^{-1} \pm 3.15 \times 10^{-2} \text{ Bq/g}$ ($6.27 \times 10^0 \pm 8.51 \times 10^{-1} \text{ pCi/g}$) at SEC to $5.11 \times 10^{-1} \pm 6.73 \times 10^{-2} \text{ Bq/g}$ ($1.38 \times 10^1 \pm 1.82 \times 10^0 \text{ pCi/g}$) at SMR. Potassium-40 concentrations in deep soils were lowest at WFF ($2.09 \times 10^{-1} \pm 2.76 \times 10^{-2} \text{ Bq/g}$ ($5.65 \times 10^0 \pm 7.46 \times 10^{-1} \text{ pCi/g}$)) and highest at SMR ($5.62 \times 10^{-1} \pm 7.29 \times 10^{-2} \text{ Bq/g}$ ($1.52 \times 10^1 \pm 1.97 \times 10^0 \text{ pCi/g}$)).

The concentration of ^{40}K was not significantly different between depths or between years. The range of concentrations observed is consistent with the average natural ^{40}K concentration in soils around the world ($4.00 \times 10^{-1} \text{ Bq/g}$ [$1.08 \times 10^1 \text{ pCi/g}$]; NCRP, 1994).

Cesium-137 was detected in 17 of the 18 soil samples (Table 4.13). In surface soils, concentrations ranged from $9.29 \times 10^{-4} \pm 5.14 \times 10^{-4} \text{ Bq/g}$ ($2.51 \times 10^{-2} \pm 1.39 \times 10^{-2} \text{ pCi/g}$) at SMR to $8.88 \times 10^{-3} \pm 1.22 \times 10^{-3} \text{ Bq/g}$ ($2.40 \times 10^{-1} \pm 3.30 \times 10^{-2} \text{ pCi/g}$) at MLR. The concentration in intermediate depth soils ranged from $1.16 \times 10^{-3} \pm 4.33 \times 10^{-4} \text{ Bq/g}$ ($3.14 \times 10^{-2} \pm 1.17 \times 10^{-2} \text{ pCi/g}$) at SMR to $1.65 \times 10^{-2} \pm 2.16 \times 10^{-3} \text{ Bq/g}$ ($4.46 \times 10^{-1} \pm 5.84 \times 10^{-2} \text{ pCi/g}$) at MLR. In deep soils, the lowest concentrations of ^{137}Cs were found at SMR ($7.55 \times 10^{-4} \pm 5.00 \times 10^{-4} \text{ Bq/g}$ [$2.04 \times 10^{-2} \pm 1.35 \times 10^{-2} \text{ pCi/g}$]) and the highest concentrations were found at MLR ($1.48 \times 10^{-2} \pm 2.01 \times 10^{-3} \text{ Bq/g}$ [$4.00 \times 10^{-1} \pm 5.43 \times 10^{-2} \text{ pCi/g}$]).

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

Table 4.13 - Selected Radionuclide Concentrations (Bq/g) in Soil Near the WIPP Site.
See Appendix B for the sampling locations.

Location	Depth (cm)	[RN] ^a	2 × TPU ^b	MDC ^c	[RN]	2 × TPU	MDC
		¹³⁷ Cs			⁶⁰ Co		
MLR	0-2	8.88×10 ⁻³	1.22×10 ⁻³	4.29×10 ⁻⁴	2.97×10 ⁻⁴	3.03×10 ⁻⁴	4.81×10 ⁻⁴
MLR	2-5	1.65×10 ⁻²	2.16×10 ⁻³	5.81×10 ⁻⁴	-1.72×10 ⁻⁴	4.88×10 ⁻⁴	8.25×10 ⁻⁴
MLR	5-10	1.48×10 ⁻²	2.01×10 ⁻³	8.07×10 ⁻⁴	2.20×10 ⁻⁶	4.33×10 ⁻⁴	7.33×10 ⁻⁴
SEC	0-2	2.82×10 ⁻³	5.48×10 ⁻⁴	5.14×10 ⁻⁴	2.28×10 ⁻⁴	2.34×10 ⁻⁴	3.68×10 ⁻⁴
SEC	2-5	2.15×10 ⁻³	4.51×10 ⁻⁴	4.51×10 ⁻⁴	7.22×10 ⁻⁵	2.48×10 ⁻⁴	4.22×10 ⁻⁴
SEC	5-10	9.44×10 ⁻⁴	3.77×10 ⁻⁴	5.29×10 ⁻⁴	8.55×10 ⁻⁵	2.80×10 ⁻⁴	4.74×10 ⁻⁴
SMR	0-2	9.29×10 ⁻⁴	5.14×10 ⁻⁴	7.81×10 ⁻⁴	3.66×10 ⁻⁴	4.29×10 ⁻⁴	6.92×10 ⁻⁴
SMR	2-5	1.16×10 ⁻³	4.33×10 ⁻⁴	6.03×10 ⁻⁴	1.82×10 ⁻⁴	3.20×10 ⁻⁴	5.25×10 ⁻⁴
SMR	5-10	7.55×10 ⁻⁴	5.00×10 ⁻⁴	7.73×10 ⁻⁴	2.11×10 ⁻⁴	4.63×10 ⁻⁴	7.59×10 ⁻⁴
WEE	0-2	2.21×10 ⁻³	5.48×10 ⁻⁴	6.55×10 ⁻⁴	8.29×10 ⁻⁵	2.48×10 ⁻⁴	4.22×10 ⁻⁴
WEE	2-5	3.13×10 ⁻³	5.33×10 ⁻⁴	5.62×10 ⁻⁴	-3.74×10 ⁻⁴	4.14×10 ⁻⁴	7.10×10 ⁻⁴
WEE	5-10	2.49×10 ⁻³	6.22×10 ⁻⁴	7.62×10 ⁻⁴	-5.03×10 ⁻⁴	4.63×10 ⁻⁴	7.88×10 ⁻⁴
WFF	0-2	3.57×10 ⁻³	5.40×10 ⁻⁴	3.59×10 ⁻⁴	-6.99×10 ⁻⁵	1.75×10 ⁻⁴	3.40×10 ⁻⁴
WFF	2-5	4.48×10 ⁻³	6.22×10 ⁻⁴	3.89×10 ⁻⁴	-1.96×10 ⁻⁵	2.76×10 ⁻⁴	4.66×10 ⁻⁴
WFF	5-10	3.30×10 ⁻³	4.88×10 ⁻⁴	2.63×10 ⁻⁴	2.13×10 ⁻⁴	1.61×10 ⁻⁴	2.46×10 ⁻⁴
WSS	0-2	1.95×10 ⁻³	4.11×10 ⁻⁴	3.96×10 ⁻⁴	5.85×10 ⁻⁵	2.42×10 ⁻⁴	4.14×10 ⁻⁴
WSS	2-5	1.78×10 ⁻³	5.07×10 ⁻⁴	6.51×10 ⁻⁴	-3.08×10 ⁻⁴	4.22×10 ⁻⁴	7.18×10 ⁻⁴
WSS	5-10	1.10×10 ⁻³	3.29×10 ⁻⁴	4.07×10 ⁻⁴	2.24×10 ⁻⁴	2.49×10 ⁻⁴	3.96×10 ⁻⁴
		⁹⁰ Sr			⁴⁰ K		
MLR	0-2	-2.81×10 ⁻³	1.14×10 ⁻²	1.89×10 ⁻²	4.51×10 ⁻¹	5.96×10 ⁻²	9.14×10 ⁻³
MLR	2-5	9.62×10 ⁻³	1.27×10 ⁻²	2.00×10 ⁻²	4.77×10 ⁻¹	6.25×10 ⁻²	1.38×10 ⁻²
MLR	5-10	-3.96×10 ⁻⁴	1.04×10 ⁻²	1.72×10 ⁻²	4.29×10 ⁻¹	5.66×10 ⁻²	1.37×10 ⁻²
SEC	0-2	4.37×10 ⁻³	1.21×10 ⁻²	2.05×10 ⁻²	2.24×10 ⁻¹	3.05×10 ⁻²	5.62×10 ⁻³
SEC	2-5	1.29×10 ⁻²	1.21×10 ⁻²	1.97×10 ⁻²	2.32×10 ⁻¹	3.15×10 ⁻²	5.85×10 ⁻³
SEC	5-10	1.17×10 ⁻²	1.33×10 ⁻²	2.18×10 ⁻²	2.51×10 ⁻¹	3.39×10 ⁻²	5.48×10 ⁻³
SMR	0-2	-1.34×10 ⁻³	1.26×10 ⁻²	2.07×10 ⁻²	4.74×10 ⁻¹	6.18×10 ⁻²	1.34×10 ⁻²
SMR	2-5	-2.93×10 ⁻³	1.17×10 ⁻²	1.94×10 ⁻²	5.11×10 ⁻¹	6.73×10 ⁻²	9.47×10 ⁻³
SMR	5-10	3.37×10 ⁻³	1.25×10 ⁻²	2.13×10 ⁻²	5.62×10 ⁻¹	7.29×10 ⁻²	1.37×10 ⁻²
WEE	0-2	1.79×10 ⁻²	9.55×10 ⁻³	1.60×10 ⁻²	2.25×10 ⁻¹	3.07×10 ⁻²	7.36×10 ⁻³
WEE	2-5	-8.70×10 ⁻³	9.07×10 ⁻³	1.60×10 ⁻²	2.50×10 ⁻¹	3.41×10 ⁻²	1.33×10 ⁻²
WEE	5-10	-3.15×10 ⁻³	9.47×10 ⁻³	1.63×10 ⁻²	2.62×10 ⁻¹	3.55×10 ⁻²	1.32×10 ⁻²
WFF	0-2	-3.40×10 ⁻³	9.32×10 ⁻³	1.61×10 ⁻²	2.53×10 ⁻¹	3.31×10 ⁻²	4.33×10 ⁻³
WFF	2-5	-4.22×10 ⁻³	9.14×10 ⁻³	1.58×10 ⁻²	2.58×10 ⁻¹	3.40×10 ⁻²	1.06×10 ⁻²
WFF	5-10	-4.22×10 ⁻³	9.51×10 ⁻³	1.65×10 ⁻²	2.09×10 ⁻¹	2.76×10 ⁻²	4.63×10 ⁻³
WSS	0-2	9.62×10 ⁻³	1.25×10 ⁻²	2.07×10 ⁻²	2.29×10 ⁻¹	3.12×10 ⁻²	6.96×10 ⁻³
WSS	2-5	7.92×10 ⁻³	1.24×10 ⁻²	2.06×10 ⁻²	2.44×10 ⁻¹	3.34×10 ⁻²	1.33×10 ⁻²
WSS	5-10	4.07×10 ⁻³	1.20×10 ⁻²	2.03×10 ⁻²	2.35×10 ⁻¹	3.20×10 ⁻²	5.74×10 ⁻³

^a [RN] = Radionuclide concentration

^b Total propagated uncertainty

^c Minimum detectable concentration

Although ¹³⁷Cs is a fission product, it is ubiquitous in soils because of global fallout from atmospheric nuclear weapons testing. In 1998, prior to WIPP accepting any waste, the average concentration of ¹³⁷Cs in soils around WIPP was 4.3×10⁻³ Bq/g (1.16×10⁻¹ pCi/g). There was no statistically significant difference between concentrations measured in 2000 and 2001 (ANOVA, p = 0.519).

Waste Isolation Pilot Plant 2001 Site Environmental Report
DOE/WIPP 02-2225

Strontium-90 was only detected at WEE, depth 0-2cm, slightly above the MDC (Table 4.13). However, ^{60}Co was not detected in any of the soil samples. There was nonsignificant difference in ^{60}Co concentrations between years ($p = 0.421$).

Soil samples collected from one location (WEE) were divided into two parts and analyzed separately (Table 4.14). Uranium-234, ^{238}U , ^{40}K , and ^{137}Cs were compared between the duplicates. Other radionuclides of interest had insufficient detections to allow a reasonable comparison. The RER was greater than one for ^{234}U and ^{238}U in all samples. However, it was less than one in one for ^{40}K analyses and ^{137}Cs analyses. A paired t-test indicated no significant difference between ^{234}U duplicates ($p = 0.624$) and between ^{238}U duplicates ($p = 0.666$). This circumstance indicates a lack of precision in these analyses, primarily due to the non-homogeneous distribution of radionuclides in soils. Because of small-scale differences in topography, soil type and structure, soil moisture, and other microenvironmental conditions, radionuclides are rarely homogeneously distributed in soils, and good agreement between duplicate samples is difficult to achieve. However, all the measurements were low, within the range of natural concentrations, and did not differ in time or space in such a way as to suggest WIPP-related contamination of the environment.

Table 4.14 - Results of Duplicate Soil Sample Analysis. Units are Bq/g.
See Appendix B for the sampling locations.

Location	Depth (cm)	[RN] ^a	2×TPU ^b	MDC ^c	RER ^d	[RN]	2×TPU ^a	MDC ^b	RER ^c
		^{234}U				^{238}U			
WEE	0-2	5.66×10^{-3}	1.35×10^{-3}	1.02×10^{-4}	2.94	6.85×10^{-3}	1.57×10^{-3}	1.01×10^{-4}	2.26
WEE D ^e	0-2	1.50×10^{-2}	2.88×10^{-3}	8.92×10^{-4}		1.39×10^{-2}	2.70×10^{-3}	8.88×10^{-5}	
WEE	2-5	6.73×10^{-3}	1.38×10^{-3}	7.81×10^{-5}	2.63	7.33×10^{-3}	1.48×10^{-3}	7.77×10^{-5}	2.81
WEE D ^e	2-5	1.57×10^{-2}	3.12×10^{-3}	9.99×10^{-5}		1.81×10^{-2}	3.53×10^{-3}	9.92×10^{-5}	
WEE	5-10	1.84×10^{-2}	3.68×10^{-3}	2.83×10^{-4}	1.94	2.00×10^{-2}	3.96×10^{-3}	1.04×10^{-4}	1.90
WEE D ^e	5-10	1.01×10^{-2}	2.20×10^{-3}	3.00×10^{-4}		1.12×10^{-2}	2.39×10^{-3}	3.77×10^{-4}	
		^{40}K				^{137}Cs			
		[RN]	2×TPU	MDC	RER	[RN]	2×TPU	MDC	RER
WEE	0-2	2.25×10^{-1}	3.07×10^{-2}	7.36×10^{-3}	0.40	2.21×10^{-3}	5.48×10^{-4}	6.55×10^{-4}	0.25
WEE D ^e	0-2	2.43×10^{-1}	3.30×10^{-2}	6.29×10^{-3}		2.40×10^{-3}	5.22×10^{-4}	5.55×10^{-4}	
WEE	2-5	2.62×10^{-1}	3.55×10^{-2}	1.32×10^{-2}	0.41	2.49×10^{-3}	6.22×10^{-4}	7.62×10^{-4}	0.18
WEE D ^e	2-5	2.42×10^{-1}	3.28×10^{-2}	6.36×10^{-3}		2.35×10^{-3}	4.74×10^{-4}	4.55×10^{-4}	
WEE	5-10	2.62×10^{-1}	3.55×10^{-2}	1.32×10^{-2}	0.22	2.49×10^{-3}	6.22×10^{-4}	7.62×10^{-4}	0.50
WEE D ^e	5-10	2.51×10^{-1}	3.42×10^{-2}	1.34×10^{-2}		2.12×10^{-3}	4.07×10^{-4}	5.14×10^{-4}	

^a [RN] = Radionuclide concentration

^b Total propagated uncertainty

^c Minimum detectable concentration

^d Relative Error Ratio

^e Duplicate

4.7 Sediments

4.7.1 Sample Collection

Sediment samples were collected from 12 locations around the WIPP site, mostly from the same water bodies from which the surface water samples were collected (Figure 4.7, see Appendix B for location codes). The samples were collected in 1 l